

# City of Rancho Palos Verdes GHG Inventory, Forecasting, Target-Setting Report for an Energy Efficiency Climate Action Plan

# January 2015

Prepared for:



Prepared by: ATKINS 3570 Carmel Mountain Road, Suite 300 San Diego, California 92130

Funded by:



Local Government Strategic Plan Strategies Program Under the auspices of the California Public Utilities Commission

## Contents

List of Acronyms and Abbreviations iii
Key Findings iv
Introduction1
GHG Emissions Inventories 2
Emissions Reporting
Community Emissions
Energy
2005—2012 Emissions Summary
Inventory Forecasts
Business-as-Usual Forecasts       17         Community Business-as-Usual Forecast       17         Municipal Business-as-Usual Forecast       19         Adjusted Business-as-Usual Forecasts       19         Community Adjusted Business-as-Usual Forecast       20         Municipal Adjusted Business-as-Usual Forecast       21
Reduction Targets
Recommended Community Targets
Conclusions and Next Steps
References

Appendix A: Glossary of Terms Appendix B: Methodology

#### Tables

Table 1. Key Terms in the Report	1
Table 2. GHGs Analyzed in the Inventories	
Table 3. Community-Wide GHG Emissions by Sector for 2005 and 2012	6
Table 4. Community GHG Emissions for 2005, 2007, 2010, and 2012	7
Table 5. Activity Data used in 2005, 2007, 2010, and 2012 Community Inventories	8
Table 6. Demographic Data for 2005, 2007, 2010, and 2012	9
Table 7. Activity Data and GHG Emissions of Energy in 2005 and 2012	9
Table 8. Municipal GHG Emissions by Sector for 2005 and 2012	12
Table 9. Municipal GHG Emissions for 2005, 2007, 2010, and 2012	
Table 10. Activity Data used in 2005, 2007, 2010, and 2012 Municipal Inventories	14
Table 11. Activity Data and GHG Emissions of Energy in 2005 and 2012	15
Table 12. Growth Factors for 2012, 2020, and 2035	18
Table 13. Community BAU Forecast	18
Table 14. Municipal BAU Forecast	19
Table 15. Community Adjusted BAU Emissions	
Table 16. Municipal Adjusted BAU Emissions	
Table 17. State-Aligned Community GHG Reduction Targets	
Table 18. State-Aligned Municipal GHG Reduction Targets	23

#### Figures

Figure 1. Community-Wide GHG Emissions by Sector for 2005 and 2012	5
Figure 2. Community GHG Emissions for 2005, 2007, 2010, and 2012	6
Figure 3. GHG Emissions for Community Electricity and Natural Gas, by Sector	10
Figure 4. Municipal GHG Emissions by Sector for 2005 and 2012	12
Figure 5. Municipal GHG Emissions for 2005, 2007, 2010, and 2012	13
Figure 6. GHG Emissions for Municipal Electricity and Natural Gas, by Sector	16
Figure 7. Community Emissions Inventories, Projections, and Targets	23
Figure 8. Municipal Emissions Inventories, Projections, and Targets	24

# List of Acronyms and Abbreviations

AB	Assembly Bill
ADC	Alternative Daily Cover
BAU	Business-as-Usual
CAFE	Corporate Average Fuel Economy
CH <sub>4</sub>	Methane
CARB	California Air Resources Board
CIWMB	California Integrated Waste Management Board
CO <sub>2</sub>	Carbon Dioxide
CO <sub>2</sub> e	Carbon Dioxide Equivalents
EECAP	Energy Efficiency Climate Action Plan
EO	Executive Order
GHG	Greenhouse Gas
GWP	Global Warming Potential
IEAP	International Local Government GHG Emissions Analysis Protocol
IFT	Inventories, Long-Term Forecasts, and Target-Setting
IPCC	Intergovernmental Panel on Climate Change
JWPCP	Joint Water Pollution Control Plant
kWh	Kilowatt-hour
LCFS	Low Carbon Fuel Standard
LGOP	Local Government Operations Protocol
MT	Metric Tons
NDN	Nitrification/denitrification
N <sub>2</sub> O	Nitrous Oxide
RPS	Renewable Portfolio Standard
RTP	Regional Transportation Plan
SBCCOG	South Bay Cities Council of Governments
SCAG	Southern California Association of Governments
SCE	Southern California Edison
SCG	Southern California Gas Company
SEEC	Statewide Energy Efficiency Collaborative

### **Key Findings**

#### Community

- The City of Rancho Palos Verdes decreased emissions 8% from 2005 to 2012, from 287,025 MT CO<sub>2</sub>e to 263,876 MT CO<sub>2</sub>e.
- On-Road Transportation, Residential Energy, Solid Waste, Water, and Wastewater sector emissions decreased while Commercial Energy and Off-road Sources sectors increased emissions from 2005 to 2012.
- Energy-related emissions accounted for 38% of the total community emissions in 2005 and 42% of total emissions in 2012.
- Under the Adjusted Business-as-Usual (BAU) scenario, emissions will be 229,481 MT  $CO_2e$  in 2020 and 190,653 MT  $CO_2e$  in 2035. These emissions levels are 20% lower in 2020 than 2005 levels and 34% lower than 2005 levels by 2035.
- The City should choose a reduction target that is feasible and ambitious. The State recommends a 15% reduction below 2005 levels by 2020, which would be achieved under the Adjusted BAU scenario.
- To continue reductions consistent with the State's long-term emissions reduction goal of lowering emissions 80% below 1990 levels by 2050, the City would need to reduce emissions in 2035 by 44,270 MT CO<sub>2</sub>e from an Adjusted BAU forecast. This is a 23% reduction from the 2035 Adjusted BAU emissions level and would achieve a 49% reduction from 2005 levels.

#### Municipal

- Municipal emissions have increased approximately 1% from 2005 to 2012, from 2,264 MT  $CO_2e$  to 2,291 MT  $CO_2e$ .
- Emissions in the Employee Commute, City-owned Outdoor Lights, Vehicle Fleet & Equipment, and Solid Waste decreased during the period.
- Municipal emissions are a subset of community emissions and account for less than 1% of all emissions.
- Under the Adjusted BAU scenario, emissions will be 2,177 MT  $CO_2e$  in both 2020 and 2035. These emissions levels are nearly 4% lower than in 2005.
- The City will need to reduce emissions by 253 MT CO<sub>2</sub>e from the 2020 Adjusted BAU emissions level to meet a 15% reduction target from 2005 levels. By 2035, the City will need to reduce emissions by 1,155 MT CO<sub>2</sub>e from the 2035 Adjusted BAU emissions level to meet a 49% reduction target from 2005 levels.

### Introduction

The Greenhouse Gas (GHG) Inventories, Long-Term Forecasts, and Target-Setting (IFT) Report contains the first steps toward the City of Rancho Palos Verdes (City) identifying energy-efficiency measures in an Energy Efficiency Climate Action Plan (EECAP). The inventories describe historic energy use and GHG emissions and the forecasts describe projected future emissions in the City. The target-setting section describes GHG reduction recommendations that are consistent with State goals and may assist the City in establishing local GHG reduction targets. The inventories and recommended reduction targets will help the City in the next step of the EECAP, which is to identify energy efficiency and GHG reduction measures that are relevant, meaningful, and feasible.

Specifically, the IFT Report includes (words and phrases in bold are described in Table 1):

- Historic GHG emissions in community inventories and municipal inventories for 2005, 2007, 2010, and 2012;
- Future GHG emissions for 2020 and 2035 under a **business-as-usual** forecast scenario and **adjusted business-as-usual** forecast scenario; and
- Recommended GHG reduction targets for 2020 and 2035.

Term	Definition
Adjusted business-as-usual	A GHG forecast scenario that accounts for known policies and regulations that will affect future emissions. Generally, these are state and federal initiatives that will reduce emissions from the business-as-usual scenario.
Baseline year	The inventory year used for setting targets and comparing future inventories against.
Business-as-usual	A GHG forecast scenario that assumes no change in policy affecting emissions since the most recent inventory. Changes in emissions are driven primarily through changes in demographics.
Community Inventory	GHG emissions that result from the activities by residents and businesses in the city. An inventory reports emissions that occur over a single calendar year.
Emission factors	The GHG-intensity of an activity.
Municipal Inventory	GHG emissions that result from the activities performed as part of the government operations in the city and are a subset of the community inventory. An inventory reports emissions that occur over a single calendar year.
Reduction targets	GHG emissions levels not to be exceeded by a specific date. Local reduction targets are often informed by state recommendations and different targets may be established for different years.
Sector	A subset of the emissions inventory classified by a logical grouping such as economic or municipal-specific category.

#### Table 1. Key Terms in the Report<sup>1</sup>

<sup>&</sup>lt;sup>1</sup> A glossary of terms is also included as Appendix A.

### **GHG Emissions Inventories**

GHG emissions inventories are the foundation of planning for future reductions. Establishing an existing inventory of emissions helps to identify and categorize the major sources of emissions currently being produced. In this report, four years of historic inventories are presented to show not only the major sources of emissions in the City, but also how those sources vary over time. For both the community and municipal inventories, the years 2005, 2007, 2010, and 2012 are presented. The 2005 inventory (for both community and municipal operations) is considered the **baseline year**. A baseline year is established as a starting point against which other inventories may be compared and targets may be set, and is generally the earliest year with a full emissions inventory. The most recent inventory (2012) has the most relevant data for planning purposes, while the interim years (2007 and 2010) provide context and may help identify trends or anomalies.

#### **Emissions Reporting**

The primary GHGs from the community and municipal operations are from carbon dioxide ( $CO_2$ ), methane ( $CH_4$ ), and nitrous oxide ( $N_2O$ ). Because each of these gases has a different capacity for trapping heat in the atmosphere, known as its global warming potential (GWP), a method of reporting is needed to be able to compare gases in the same terms. As a result, emissions are reported in carbon dioxide equivalents, or  $CO_2e$ , with each GHG normalized and calculated relative to  $CO_2$  using its GWP. Table 2 describes the GHGs analyzed in this report, their symbol, GWP, and primary community sources of emissions. While  $N_2O$  has the highest GWP and may be considered the most dangerous on a permolecule basis,  $CO_2$  is by far the most prevalent, accounting for 88% of statewide emissions in 2005 (CARB 2011).

Greenhouse Gas	Symbol	Global Warming Potential	Primary Community Sources
Carbon Dioxide	CO <sub>2</sub>	1	Fossil fuel combustion
Methane	CH <sub>4</sub> 25		Fossil fuel combustion, landfills, wastewater treatment
Nitrous Oxide	N <sub>2</sub> O	298	Fossil fuel combustion, wastewater treatment

Table 2. GHGs Analyzed in the Inventories

Source: IPCC Fourth Assessment Report, 2007.

#### **Emissions Sectors**

The inventories identify the major sources of GHGs emissions caused by activities in sectors that are specific to community or municipal activities. A **sector** is a subset of the economy, society, or municipal operations whose components share similar characteristics. An emissions sector can also contain subsectors that provide more specificity about the source of emissions (e.g., natural gas and electricity are subsectors of the energy sector).

As mentioned above, inventories were completed for the community and municipal operations. Because the majority of municipal activities occur within the boundaries of the City and therefore contribute to the overall emissions of the community, both inventories are interconnected, with the municipal inventory considered a subset of the community inventory. As a result, municipal emissions are included in numbers reported for the community. The municipal inventory is separated to highlight areas of emissions that the City has more direct control over and to identify where they can begin to set examples for the community on how reduction strategies can be implemented.

The following subsections describe the sectors used in the community and municipal inventories. It is important to note that both inventories capture similar types of information but may be categorized differently. For example, energy is reported in both the community and municipal inventory, but community level energy emissions are reported as "Residential" and "Commercial/Industrial", whereas municipal energy emissions are more logically reported as "Buildings & Facilities" and "Outdoor Lights."<sup>2</sup>

#### **Community Sectors**

The community inventory is categorized by sectors based on the sector's ability to be affected through regional and local programs, incentives, zoning, and other policies. The City's community inventories were divided into the following sectors:

- **Energy** in the Community Inventory is further broken down into two sectors:
  - **Commercial/Industrial Energy** includes emissions from electricity and natural gas consumption in non-residential buildings and facilities (including outdoor lights) in the City.
  - **Residential Energy** includes emissions from electricity and natural gas consumption in residential buildings in the City.
- **On-road Transportation** includes emissions from vehicle fuel use in trips wholly within the City (in-boundary) and trips that either originate or end in the City (cross-boundary). Emissions from in-boundary trips are fully accounted for in the inventory, whereas only half of the emissions from cross-boundary trips are accounted for. Trips that pass-through the City (such as CA-213,) are not accounted for in the inventory because the City has little or no control of these emissions. As a result, this methodology reflects only trips or parts of trips within City borders that the City has the ability to affect.
- **Solid Waste** includes emissions from waste that is generated in the community and sent to landfills.
- **Water** includes emissions from the electricity used to source, treat, and deliver imported water in the community that is not accounted for in the community utility data.
- Wastewater includes emissions from treating wastewater generated in the community.
- **Off-road Sources** include emissions from operating equipment for construction, commercial, light industrial and agricultural activities; lawn and garden equipment; and recreational vehicles such as all-terrain vehicles.

<sup>&</sup>lt;sup>2</sup> Outdoor Lights are further categorized as SCE-owned Outdoor Lights or City-owned Outdoor Lights as described later.

#### **Municipal Sectors**

Sources of municipal emissions are divided into the following sectors:

- Energy in the municipal inventory is further broken down into four sectors:
  - **Buildings and Facilities** includes energy use by the government, including electricity and natural gas.
  - SCE-owned Outdoor Lights includes electricity for streetlights on fixtures owned by SCE.
  - **City-owned Outdoor Lights** includes electricity for streetlights on fixtures owned by the City and traffic control signals.
  - **Water Delivery** includes electricity for water and sewer pumping, dewatering wells, and irrigation.
- Vehicle Fleet & Equipment includes emissions from vehicles owned or operated by the government or contracted by the City for services such as street cleaning. It also includes equipment, such as emergency generators.
- **Employee Commute** includes emissions from fuel use in vehicle trips by municipal employees commuting to and from work in the City.
- **Solid Waste** includes emissions from waste generated by municipal employees or at municipally-owned facilities.

#### **Calculation Methodology**

GHG emissions were calculated using activity data available (e.g., kilowatt-hours of electricity) for each sector and protocols for converting activity data to emissions output using relevant **emission factors**. Emission factors relate the activity to GHG emissions and may vary by year (e.g., for electricity) and often are not affected by local actions or behavior, unlike activity data. The U.S. Community Protocol for Accounting and Reporting Greenhouse Gas Emissions (ICLEI 2012) and the Local Government Operations Protocol for the Quantification and Reporting of GHG Emissions Inventories (LGOP) (CARB 2010) were the primary protocols used for developing the community and municipal inventories, respectively. Activity data are reported in the community and municipal emissions subsections below, and emission factors are detailed in Appendix B.

#### **Community Emissions**

The community inventory includes the GHG emissions that result from activities within City boundaries. This section presents the findings of the community inventory for four years: 2005 (baseline year), 2007, 2010, and 2012. It also provides more specific detail and findings on the energy sectors, which will form the basis of the reduction targets and reduction measures the City identifies in the EECAP.

#### 2005—2012 Emissions Summary

The City of Rancho Palos Verdes reduced emissions 8% from 2005 to 2012, from 287,025 MT CO<sub>2</sub>e to 263,876 MT CO<sub>2</sub>e.

 On-road Transportation, Residential Energy, Solid Waste, Water, and Wastewater sector emissions decreased while the Commercial Energy and Off-road Sources sectors increased emissions from 2005 to 2012.

As shown in Figure 1 and Table 3, the Transportation sector was the largest contributor to emissions in both 2005 (53%) and 2012 (52%) by producing 150,564 MT CO<sub>2</sub>e in 2005 and 136,175 MT CO<sub>2</sub>e in 2012. This change represents almost 10% decrease in emissions from 2005 to 2012. Residential energy is the second-largest contributor to emissions, adding 31% in 2005 and 33% 2012. The total Residential energy emissions decreased by about 3% from 2005 to 2012, from 88,941 MT CO<sub>2</sub>e to 86,129 MT CO<sub>2</sub>e. Commercial Energy represented 7% of emissions in 2005 and 10% in 2012, and its total emissions increased by about 24%, from 20,377 MT CO<sub>2</sub>e in 2005 to 25,304 MT CO<sub>2</sub>e in 2012. Water comprised 6% of the total (18,156 MT CO<sub>2</sub>e) in 2005, but was reduced to 4% of the total (11,653 MT CO<sub>2</sub>e) in 2012. Solid Waste, Wastewater, and Off-road Sources made up the remaining emissions in each year. Solid Waste and Wastewater emissions declined from 2005 to 2012; however, Off-road Sources increased 117% (from 157 to 340 MT CO<sub>2</sub>e) in the same period. Off-road Sources comprise a very small percentage of overall emissions, but are variable primarily due to construction-related emissions, which are based on the level of development estimated in the City each year.




Figure 1. Community-Wide GHG Emissions by Sector for 2005 and 2012

Sector	2005 (MT CO₂e)	2012 (MT CO₂e)	% Change 2005 to 2012
On-Road Transportation	150,564	136,175	-9.6%
Residential Energy	88,941	86,129	-3.2%
Commercial Energy	20,377	25,304	24.2%
Water	18,156	11,653	-35.8%
Solid Waste	8,674	4,158	-52.1%
Off-road Sources	157	340	116.7%
Wastewater	156	117	-25.0%
Total	287,025	263,876	-8.1%

#### Table 3. Community-Wide GHG Emissions by Sector for 2005 and 2012

#### 2005, 2007, 2010, and 2012 Inventories

Figure 2 and Table 4 show the GHG emissions by sector for all inventory years. Emissions are variable among the inventory years, and may reflect changes in the economy, weather, and programs implemented to reduce emissions. The table also lists the percentage of each sector relative to total emissions and shows that the proportion of each sector does not vary greatly by year.




Figure 2. Community GHG Emissions for 2005, 2007, 2010, and 2012

Sector	2005 (MT CO <sub>2</sub> e)	% of Total	2007 (MT CO <sub>2</sub> e)	% of Total	2010 (MT CO <sub>2</sub> e)	% of Total	2012 (MT CO <sub>2</sub> e)	% of Total
On-Road Transportation	150,564	52%	145,899	52%	138,685	52%	136,175	52%
Residential Energy	88,941	31%	90,581	32%	86,855	32%	86,129	33%
Commercial Energy	20,377	7%	20,209	7%	22,872	9%	25,304	10%
Water	18,156	6%	17,604	6%	11,194	4%	11,653	4%
Solid Waste	8,674	3%	8,138	3%	6,296	2%	4,158	2%
Off-road Sources	157	<1%	295	<1%	1,520	1%	340	<1%
Wastewater	156	<1%	118	<1%	117	<1%	117	<1%
Total	287,025		282,844		267,539		263,876	
% Change from 2005			-1.5%		-6.8%		-8.1%	

Table 4. Community GHG Emissions for 2005, 2007, 2010, and 2012

Activity data can provide more insight into behavioral changes in the community, as these data are not affected by emission factors. Table 5 summarizes activity data for each sector and subsector. The activity data show that Commercial/Industrial Electricity, and some Off-road Sources increased from 2005 to 2012, while Vehicle Miles Traveled, Residential Natural Gas, Solid Waste, and Water and Wastewater consumption decreased from 2005 to 2012. Residential Electricity increased less than 1%. Wastewater and Off-road Source emissions use indicator data to attribute county-level emissions to the City and the indicator data are also shown in Table 5. Notably, while On-road Transportation emissions decreased nearly 10% between 2005 and 2012, vehicle miles traveled increased by only 4%. The difference reflects that for each vehicle mile traveled, fewer emissions are generated due to improvements in the fuel efficiency of vehicles.

Demographic data also help provide perspective to changes in emissions over time. Table 6 shows the number of households, jobs, population, and service population (jobs + population) for each inventory year. Energy emissions in particular often reflect trends in demographic data. For example, decrease in Residential energy usage between 2005 and 2012 may reflect the small decline in population over the same period.

#### Table 5. Activity Data used in 2005, 2007, 2010, and 2012 Community Inventories

Sector	2005	2007	2010	2012	% Change 2005 to 2012	
On-road Transportation						
Total Vehicle Miles Traveled	287,926,109	281,650,326	274,693,126	276,102,859	-4.1%	
Residential Energy						
Electricity (kWh)	126,988,678	128,757,878	126,318,979	127,691,251	0.6%	
Natural Gas (therms)	9,471,187	10,064,521	9,495,769	8,518,368	-10.1%	
Commercial/Industrial Energy					-	
Electricity (kWh)	54,342,771	57,319,790	62,520,704	64,563,386	18.8%	
Natural Gas (therms)	727,360	697,781	917,031	876,032	20.4%	
Solid Waste						
Landfilled (tons)	33,150	31,592	25,285	16,815	-49.3%	
ADC (tons) <sup>1</sup>	3,168	1,855	382	112	-96.5%	
Water and Wastewater						
Water (Million Gallons)	3,424	3,424	3,040	3,165	-7.6%	
Wastewater (City portion of countywide residents)	0.43%	0.43%	0.42%	0.42%	-2.1%	
Off-road Sources <sup>2</sup> (% of LA County	emissions attril	outed to the City	r)		<u>.</u>	
Lawn & Garden (% Households)	0.49%	0.48%	0.47%	0.47%	-3.2%	
Construction (% Building permits)	0.05%	0.09%	0.51%	0.10%	114.4%	
Industrial (% Manufacturing jobs)	0.03%	0.03%	0.03%	0.03%	5.0%	
Light Commercial (% Other jobs)	0.15%	0.16%	0.16%	0.16%	7.8%	
Recreation (Population weighted by income)	0.93%	0.90%	0.87%	0.85%	-8.5%	
Agriculture (% Ag. Jobs)	0.00%	0.00%	0.00%	0.02%		

1: ADC is Alternative Daily Cover, which is green waste (grass, leaves, and branches) that is used to cover landfill emissions. They are reported separately by CalRecycle and therefore shown separately here.

2: Off-road Source emissions are available at the county level through CARB's OFFROAD model. Emissions attributable to the City were derived using indicator data related to the off-road source. For example, the percentage of households in the City compared to the county was used to attribute the same percentage of lawn & garden equipment emissions to the City. See Appendix B for more methodology details.

	2005	2007	2010	2012	% Change 2005-2012
Population	42,457	41,728	41,654	41,897	-1.3%
Households	15,417	15,480	15,561	15,599	1.2%
Jobs	6,191	6,575	6,132	6,298	1.7%
Service Population (Population + Jobs)	48,648	48,303	47,786	48,195	-0.9%

#### Table 6. Demographic Data for 2005, 2007, 2010, and 2012

Source: SCAG.

#### Energy

The EECAP ultimately will focus on increasing energy efficiency and reducing GHG gases from energy; therefore, it is important for the City to understand its current energy consumption to make informed decisions for reducing energy-related emissions. Energy use consists of electricity and natural gas. Emissions from Commercial/Industrial and Residential energy use account for 38% of the total community emissions in 2005 and 42% in 2012. Table 7 shows the breakdown in activity (kWh or therms) and GHG emissions by sector and energy source.

	2005		2012	2012 % Change		% Change in	
Sector	Activity (kWh or therms)	Emissions (MT CO <sub>2</sub> e)			Activity 2005-2012	Emissions 2005-2012	
Commercial/ Indus	trial						
Electricity	54,342,771	16,509	64,563,386	20,646	18.8%	25.1%	
Natural Gas	727,360	3,868	876,032	4,658	20.4%	20.4%	
Residential	Residential						
Electricity	126,988,678	38,578	127,691,251	40,833	0.6%	5.8%	
Natural Gas	9,471,187	50,363	8,518,368	45,296	-10.1%	-10.1%	
Total (MT CO₂e)		109,318		111,433		1.9%	

Table 7. Activity Data and GHG Emissions of Energy in 2005 and 2012

Commercial electricity use increased nearly 19% between 2005 and 2012; however, emissions increased by more than 25%. Similarly, Residential electricity use increased by less than 1% but emissions increased by nearly 6%. The difference between the change in activity data and emissions data are due to the emission factor used for electricity for 2005 and 2012. Emission factors convert activity data into GHG emissions and electricity emission factors vary annually based on how electricity is generated by the electricity provider (i.e., the amount of renewables, natural gas, coal, etc.). In 2005, Southern California Edison (SCE) generated electricity that resulted in an emission factor of 669.7 CO<sub>2</sub>e. In 2012, SCE's

# Electricity-Related Emissions

All emissions are comprised of activity data and the emission factor, or GHG-intensity, of that activity. For electricity, the activity data are the kilowatt-hours (kWh) used by the city's residents and businesses and the energy intensity is based on the sources of power that Southern California Edison uses to generate electricity. Changes to either component can affect the GHG emissions from electricity in the City.

electricity generation resulted in an emission factor of 705.0 CO<sub>2</sub>e. Therefore, a kilowatt-hour of electricity used in 2012 emitted more GHGs than a kilowatt-hour of electricity used in 2005. Future emissions could increase or decrease based on changes to SCE's emission factors, which the City cannot directly affect, or through changes in usage, which can be affected by changes in local policy, outreach, or incentive programs.

Unlike electricity, the emission factor for natural gas is estimated on a national basis and remains fairly constant over time. Therefore, the natural gas GHG emissions follow the same trend as usage. In Rancho Palos, Commercial/Industrial natural gas consumption (therms) increased by 20.4% from 2005 to 2012; therefore the emissions also increased 20.4%. Residential natural gas therms used and GHG emissions declined about 10% from 2005 to 2012. Figure 3 shows the trend in electricity and natural gas emissions from 2005 to 2012 for the Commercial/Industrial and Residential sectors.




Figure 3. GHG Emissions for Community Electricity and Natural Gas, by Sector

#### **Municipal Emissions**

As described earlier, a municipal GHG emissions inventory is a subset of the community inventory. The municipal inventory includes emissions from activities conducted as part of government operations in the City. While emissions from government operations are normally a fraction of the overall community emissions, the City has the most direct control over municipal emissions and the City can demonstrate leadership in the community by adopting and implementing energy and GHG reduction strategies. This section presents the findings of the municipal inventory for 2005 (the baseline year), 2007, 2010, and 2012. It also provides more specific detail and findings on the energy sectors, which will form the basis of the reduction measures the City identifies in the EECAP.

#### 2005—2012 Emissions Summary

- Municipal emissions have increased about 1% from 2005 to 2012, from 2,264 MT  $CO_2e$  to 2,291 MT  $CO_2e$ .
- Emissions decreased in the Employee Commute, City-owned Outdoor Lights, Vehicle Fleet & Equipment, and Solid Waste sectors between 2005 and 2012.
- The Vehicle Fleet & Equipment sector showed the greatest overall reductions (16 MT CO<sub>2</sub>e reductions).
- Emissions from municipal operations account for less than 1% of community emissions.

The City's Vehicle Fleet & Equipment is the sector with the largest percentage of emissions in 2005 (65%) and 2012 (64%), although emissions from this sector decreased 1% over the period (Figure 4). The second largest-emitting sector for 2005 was SCE-owned Outdoor Lights. While the percentage of total emissions from this sector remained constant between the two inventory years, at 16%, emissions increased by 4% (16 MT CO<sub>2</sub>e). The Employee Commute sector contributed 157 MT CO<sub>2</sub>e (7% of total emissions) in 2005 and decreased by 1 MT CO<sub>2</sub>e in 2012 (to 156 MT CO<sub>2</sub>e, remaining at 7% of total emissions). Emissions from the Buildings & Facilities accounted for 9% of total emissions for 2005 and 2012. Its emissions increased 10% (from 196 MT CO<sub>2</sub>e in 2005 to 215 MT CO<sub>2</sub>e in 2012). Solid Waste sector decreased emissions by 25%, from 2005 to 2012. City-owned Outdoor Lights also decreased its emissions, dropping from 35 MT CO<sub>2</sub>e to 33 MT CO<sub>2</sub>e (a 6% decrease). The Water Delivery sector increased its emissions by 42%. The 2005 and 2012 emissions and changes are detailed in Table 8.




Figure 4. Municipal GHG Emissions by Sector for 2005 and 2012

Sector	2005 (MT CO₂e)	2012 (MT CO₂e)	% Change 2005 to 2012
Fleet & Equipment	1,480	1,464	-1%
Outdoor Lights—SCE-owned	357	373	4%
Buildings & Facilities	196	215	10%
Employee Commute	157	156	-1%
Outdoor Lights—City-owned	35	33	-6%
Water Delivery	31	44	42%
Solid Waste	8	6	-25%
Total	2,264	2,291	1.2%

Table 8.	Municipal	GHG	Emissions	by Sector	for	2005	and 2012

Note: City-Owned Outdoor Lights includes streetlights and traffic signals. SCE-Owned Outdoor Lights includes streetlights on SCE-owned fixtures. Water Delivery includes water and sewer pumping, dewatering wells, and irrigation.

#### 2005, 2007, 2010, and 2012 Inventories

Figure 5 and Table 9 show the municipal GHG emissions by sector for all four inventory years. Emissions peaked in 2012 (2,291 MT  $CO_2e$ ) and were the lowest in 2010(1,991MT  $CO_2e$ ).




Figure 5. Municipal GHG Emissions for 2005, 2007, 2010, and 2012

Sector	2005 (MT CO <sub>2</sub> e)	% of Total	2007 (MT CO <sub>2</sub> e)	% of Total	2010 (MT CO <sub>2</sub> e)	% of Total	2012 (MT CO <sub>2</sub> e)	% of Total
Fleet & Equipment	1,480	65%	1,398	63%	1,323	66%	1,464	64%
Outdoor Lights— SCE-owned	357	16%	338	15%	338	17%	373	16%
<b>Buildings &amp; Facilities</b>	196	9%	235	11%	119	6%	215	9%
Employee Commute	157	7%	171	8%	156	8%	156	7%
Outdoor Lights— City-owned	35	2%	31	1%	24	1%	33	1%
Water Delivery	31	1%	35	2%	25	1%	44	2%
Solid Waste	8	<1%	8	<1%	6	<1%	6	<1%
Total	2,264		2,216		1,991		2,291	

Table 10 summarizes activity data for each sector and subsector.

Sector	2005	2007	2010	2012	% Change 2005 to 2012
Buildings & Facilities					
Electricity (kWh)	603,832	751,034	408,304	656,297	9%
Natural Gas (therms)	2,391	3,507	261	923	-61%
Outdoor Lights					
City-owned Electricity (kWh)	114,813	107,983	83,253	102,990	-10%
SCE-owned (kWh)	1,173,733	1,173,903	1,175,016	1,167,116	-1%
Fleet & Equipment					
City-owned Fleet					
Gasoline (gallons) <sup>1</sup>	3,316	2,738	2,738	2,738	-17%
Diesel (gallons)	0	0	0	0	
Contracted Fleet					
Gasoline (gallons)	24,493	26,617	10,411	16,576	-32%
Diesel (gallons) <sup>2, 3</sup>	101,226	90,259	90,259	90,259	-11%
LPG (gallons)	16,345	17,761	41,707	48,422	196%
CNG (standard cubic feet)	1,353,335	1,470,499	788,785	783,137	-42%
Employee Commute <sup>1, 3</sup>	`				<u>.</u>
Gasoline (vehicle miles traveled)	405,885	446,565	397,550	397,550	-2%
Diesel (vehicle miles traveled)	0	868	14,636	14,636	
# FTE	60	66	66	66	10%
Solid Waste <sup>1</sup>					
Generated Waste (tons)	24	26	26	26	9%
Water Delivery					
Electricity (kWh)	100,628	121,593	87,088	138,731	38%

1. Data for 2010 and 2012 were not available; therefore, activity data from 2007 data was used as a proxy.

2. ULSD gallons were used for Diesel.

3. Employee Commute survey conducted in 2014 and adjusted based on the number of employees in 2010 and 2012, which used the 2007 number of employees


#### Energy

As with the community emissions, the EECAP will focus on increasing energy efficiency and reducing GHG gases from energy within municipal operations. The City has more direct control over energyrelated emissions than other sectors, such as employee commute. Municipal energy use includes Buildings & Facilities, SCE-owned Outdoor Lights, City-owned Outdoor Lights, and Water Delivery. Energy accounted for 27% of total emissions in 2005 and 29% in 2012. While both electricity and natural gas are used for Building & Facilities, Outdoor Lights and Water Delivery only use electricity. Emissions from energy increased 7% from 2005 to 2012, with the largest overall change occurring in the Water Delivery sector (Table 11). Electricity emissions increased for the Building & Facilities, SCE-owned Outdoor Lights, and Water Delivery sectors and decreased in City-owned Outdoor Lights. As with community energy, municipal emissions use variable electricity emission factors and constant natural gas emission factors.

	2005		2012		% Change in	% Change in Emissions 2005-2012		
Sector	Activity (kWh or therms)	Emissions (MT CO <sub>2</sub> e)	Activity (kWh or therms)	Emissions (MT CO <sub>2</sub> e)	Activity 2005-2012			
Buildings & Facilities								
Electricity	603,832	183	656,297	210	9%	15%		
Natural Gas	2,391	13	923	5	-61%	-62%		
Outdoor Lights—SCE	-owned							
Electricity	1,173,733	357	1,167,116	373	-1%	4%		
Outdoor Lights—City	v-owned							
Electricity	114,813	35	102,990	33	-10%	-6%		
Water Delivery								
Electricity	100,628	31	138,731	44	38%	42%		
Total (MT CO <sub>2</sub> e)		619		665		7%		

Table 11. Activity Data and GHG Emissions of Energy in 2005 and 2012

Figure 6 shows the trend in electricity and natural gas emissions from 2005 to 2012 for the municipal energy sectors.



Note: B&F is Buildings and Facilities; NatGas is natural gas.



### **Inventory Forecasts**

GHG emissions are forecast using two scenarios: a Business-as-Usual (BAU) and an Adjusted BAU scenario. The BAU scenario describes emissions based on projected growth in population and employment and does not consider policies that will reduce emissions in the future (that is, the policies in place in 2012 are assumed to remain constant through 2035). The Adjusted BAU scenario describes emissions based on projected growth *and* considers policies that will achieve GHG reductions in the future. Policies, described in detail below, include State-adopted or approved legislation that will affect future emissions. By evaluating the two scenarios, the City can see the effect that existing policies may have on future years are forecasted for each scenario: 2020 and 2035. The 2020 forecast year is consistent with the goals identified in Assembly Bill (AB) 32, which identifies a statewide GHG reduction target by 2020. The 2035 forecast year will allow the City to develop long-term strategies to continue GHG reductions beyond 2020.

#### **Business-as-Usual Forecasts**

The BAU forecasts estimate future emissions using current (2012) consumption patterns and emission factors with the anticipated growth in the City. Anticipated growth is estimated using data from regional planning scenarios developed by the Southern California Association of Governments (SCAG), the City, and other relevant sources (Table 12). The most relevant growth factors are used to project emissions by sector. For example, future Residential Energy emissions were developed using current energy use per household (from the 2012 inventory) and the anticipated number of households in the future. Actual energy use is a function of several variables, not only the number of households; however, this approach is supported by current protocols and best practices within the State and provides a consistent approach to forecasting. Compound annual growth rates were developed using the growth projections from 2012 to 2020 and from 2021 to 2035, as shown Table 12.

In general, the City is expecting modest growth to 2035 as service population and jobs are expected to increase slightly. Although population and household are expected to slightly decrease between 2012-2020, they are expected to have no change 2020-2035. SCAG is projecting fewer vehicle miles traveled from 2012 to 2020 despite the service population and job growth. Due to the relatively low growth, the City does not anticipate major staffing changes in its government services.

#### **Community Business-as-Usual Forecast**

 BAU community emissions are expected to decrease 9% from baseline levels by 2020 and 9.5% by 2035.

The City's BAU emissions in 2020 are estimated to be 260,072 MT CO<sub>2</sub>e, or a 9% decrease from baseline (2005) emissions. By 2035, emissions are estimated to decrease 9.5% from the baseline level to 259,792 MT CO<sub>2</sub>e (Table 13).

Sector	Demographic Indicator	2012	2020	2035	2012- 2020 CAGR <sup>1</sup>	2020- 2035 CAGR <sup>1</sup>
Residential Energy	Households	15,599	15,600	15,600	0.00%	0.00%
Commercial/ Industrial Energy	Jobs	6,298	6,700	7,100	0.78%	0.39%
NA <sup>2</sup>	Population	41,897	41,700	41,700	-0.06%	0.00%
Solid Waste, Water, Wastewater, Off- road Sources	Service Population (Population + Jobs)	48,195	48,400	48,800	0.05%	0.05%
Transportation	Vehicle Miles Traveled	276,102,859	249,995,555	245,977,042	-1.23%	-0.11%
Municipal Jobs	Municipal Emissions <sup>3</sup>	66 FTE	66 FTE	66 FTE	0%	0%

#### Table 12. Growth Factors for 2012, 2020, and 2035

Source: SCAG 2012

FTE: Full-time equivalent employees.

1: Compound annual growth rate.

2: Not Applicable. Population data are shown for informational purposes but are not used for forecasting any sector.

3: The number of jobs in the City is used as an indicator for all municipal operation emissions.

Sector	2005 (MT CO <sub>2</sub> e)	2012 (MT CO <sub>2</sub> e)	2020 (MT CO <sub>2</sub> e)	% Change 2012-2020	2035 (MT CO₂e)	%Change 2012- 2035
On-Road Transportation	150,564	136,175	130,745	-4.0%	128,645	-5.5%
Residential Energy	88,941	86,129	86,134	0.0%	86,134	0.0%
Commercial Energy	20,377	25,304	26,815	6.0%	28,416	12.3%
Water	18,156	11,653	11,703	0.4%	11,799	1.3%
Solid Waste	8,674	4,158	4,176	0.4%	4,210	1.3%
Off-Road Sources	157	340	382	12.4%	469	37.9%
Wastewater	156	117	117	0.0%	119	1.7%
Total	287,025	263,876	260,072	-1.4%	259,792	-1.5%
% Change from 2005		-8.1%	-9.4%		-9.5%	

#### Table 13. Community BAU Forecast

#### **Municipal Business-as-Usual Forecast**

#### • BAU municipal emissions are expected to be 1% above baseline levels in 2020 and 2035.

The City is not anticipating growth in city services by 2020 or 2035 from current (2012) levels; therefore, the activity data for all sectors are assumed to remain constant from 2012. Therefore, the emissions in 2020 and 2035 will be equal to those in 2012 under a BAU scenario (Table 14). In 2020 and 2035, municipal emissions are estimated to be 1% above baseline emissions.

	2005 (MT CO <sub>2</sub> e)	2012 (MT CO <sub>2</sub> e)	2020 (MT CO <sub>2</sub> e)	% Change 2012-2020	2035 (MT CO <sub>2</sub> e)	% Change 2012-2035
Vehicle Fleet	1,480	1,464	1,464	0%	1,464	0%
Outdoor Lights	392	406	406	0%	406	0%
Buildings & Facilities	196	215	215	0%	215	0%
Employee Commute	157	156	156	0%	156	0%
Water Delivery	31	44	44	0%	44	0%
Solid Waste	8	6	6	0%	6	0%
Total	2,264	2,291	2,291	0%	2,291	0%
% Change from 2005		1%	1%		1%	

Table 14. M	lunicipal	BAU	Forecast
-------------	-----------	-----	----------

#### **Adjusted Business-as-Usual Forecasts**

State legislation has been approved and/or adopted that will reduce GHG emissions in the City. These policies do not require additional local action, but should be accounted for in the City's emissions forecasts to provide a more accurate picture of future emissions and the level of local action needed to reduce emissions to levels consistent with State recommendations. This forecast is called the Adjusted BAU forecast. The measures are described briefly below.

**Low Carbon Fuel Standard.** The Low Carbon Fuel Standard (LCFS) was developed as a result of Executive Order S-1-07, which mandates that the carbon intensity of transportation fuels in California are lowered 10% by 2020. The State is currently implementing this standard, which is being phased in and will achieve full implementation in 2020.

**Assembly Bill (AB) 1493 and Advanced Clean Cars.** AB 1493 directed CARB to adopt GHG standards for motor vehicles through model year 2015 that would result in reductions in GHG emissions by up to 25% in 2030. In addition, the State's Advanced Clean Cars program includes additional components that will further reduce GHG emissions statewide, including more stringent fuel efficiency standards for model years 2017—2025 and support infrastructure for the commercialization of zero-emission vehicles. CARB anticipates additional GHG reductions of 3% by 2020, 27% by 2035, and 33% by 2050<sup>3</sup>. These are also known as "Pavley I" and "Pavley II" regulations.

<sup>&</sup>lt;sup>3</sup> CARB Advanced Clean Cars Summary Sheet

**California Building Code Title 24.** California's building efficiency standards are updated regularly to incorporate new energy efficiency technologies. The code was most recently updated in 2013 and went into effect for new development in 2014. For projects implemented after January 1, 2014, the California Energy Commission estimates that the 2013 Title 24 energy efficiency standards will reduce consumption by an estimated 25% for residential buildings and 30% for commercial buildings, relative to the 2008 standards. These percentage savings relate to heating, cooling, lighting, and water heating only; therefore, these percentage savings were applied to the estimated percentage of energy use by Title 24.

**Renewable Portfolio Standard.** The Renewable Portfolio Standard (RPS) requires energy providers to derive 33% of their electricity from qualified renewable sources. This is anticipated to lower emission factors (i.e., fewer GHG emissions per kilowatt-hour used) statewide. Therefore, reductions from RPS are taken for energy embedded in water, which uses energy sources throughout the state to move from the water source area to the City. However, no credit was taken for this measure for the SCE service region (i.e., for residential and commercial electricity used in the City supplied by SCE). Analysis of SCE's current portfolio and the sources needed to replace the nuclear generation that has been taken out of service has revealed great uncertainty in how SCE's emission factors may change over time. Therefore, the emission factor used in the 2012 inventory and the BAU forecast was also used in the Adjusted BAU forecast.

**Senate Bill X7-7.** California's SB X7-7 requires water suppliers to reduce urban per capita water consumption 20% from a baseline level by 2020. The City is supplied by Golden State Water and the reductions in GHG emissions from SB X7-7 were calculated by applying the reduction goals established by Golden State Water to the City's population in 2020 and 2035.

#### **Community Adjusted Business-as-Usual Forecast**

• Emissions are expected to decrease under the Adjusted BAU forecast and will be 20% lower in 2020 than 2005 and 34% lower than 2005 levels by 2035.

The City's Adjusted BAU emissions in 2020 are estimated to be 229,481 MT  $CO_2e$  in 2020 and 190,653 MT  $CO_2e$  in 2035 (Table 15). This change represents a 20% reduction from 2005 by 2020 and 34% reduction by 2035. Due to the stringent State vehicle standards, the emissions from the Transportation sector are expected to decrease significantly over time, while the proportion of emissions from Commercial Energy will increase. Emissions from Residential Energy, and Water and Wastewater are expected to decrease over time. Emissions from Solid Waste are expected to increase slightly over time but account for about 2% of total emissions.

Sector	2005 (MT CO₂e)	2012 (MT CO <sub>2</sub> e)	2020 (MT CO <sub>2</sub> e)	2020 % of Total	2035 (MT CO <sub>2</sub> e)	2035 % of Total
Transportation & Mobile Sources	150,721	136,515	105,178	46%	65,764	34%
Residential Energy	88,941	86,129	86,134	38%	86,134	45%
Commercial Energy	20,377	25,304	26,555	12%	27,816	15%
Water & Wastewater	18,312	11,770	7,438	3%	6,729	4%
Solid Waste	8,674	4,158	4,176	2%	4,210	2%
Total	287,025	263,876	229,481	100%	190,653	100%
% Change from 2005		-8%	-20%		-34%	

#### **Municipal Adjusted Business-as-Usual Forecast**

• Under an Adjusted BAU forecast, the City's emissions will be 4% below 2005 levels in 2020 and 2035.

The City's Municipal Adjusted BAU emissions in 2020 are estimated to be 2,177 MT CO<sub>2</sub>e, which is 4% below the 2005 baseline level (Table 16). By 2035, the level of reductions is also anticipated to be 4% below 2005 levels, under the assumption that government services will neither expand nor contract significantly. The Adjusted BAU emissions are slightly lower than the BAU emissions due to the Low Carbon Fuel Standard measure described earlier. The Low Carbon Fuel Standard would lower the carbon intensity of fuels used in both the City's Vehicle Fleet and Employee Commute sectors.

Sector	2005 (MT CO <sub>2</sub> e)	2012 (MT CO <sub>2</sub> e)	2020 (MT CO <sub>2</sub> e)	2020 % of Total	2035 (MT CO <sub>2</sub> e)	2035 % of Total
Vehicle Fleet	1,480	1,464	1,361	63%	1,361	63%
Outdoor Lights	392	406	406	19%	406	19%
Buildings & Facilities	196	215	215	10%	215	10%
Employee Commute	157	156	145	7%	145	7%
Water Delivery	31	44	44	2%	44	2%
Solid Waste	8	6	6	<1%	6	<1%
Total	2,264	2,291	2,177	100%	2,177	100%
% Change from 2005		1%	-4%		-4%	

### **Reduction Targets**

The State has set goals for reducing GHG emissions by 2020 and 2050 through AB 32 and Executive Order (EO) S-3-05, respectively. The State has also provided guidance to local jurisdictions as "essential partners" in achieving the State's goals by identifying a 2020 recommended reduction goal. That goal, stated in the AB 32 Scoping Plan, was for local governments to achieve a 15% reduction below 2005 levels by 2020, which aligns with the State's goal of not exceeding 1990 emissions levels by 2020<sup>4</sup>. The State's long term target is to emit no more than 20% of 1990 levels by 2050 (or, a reduction of 80% below 1990 levels by 2050). The State has not provided an interim target, nor has it provided guidance to local governments beyond the 2020 emissions target recommendations. It is, however, clear that the issue of climate change will not end in 2020 and continued reductions should be achieved to keep the State on a path toward the 2050 goal. A straight-line projection from the 2020 to 2050 goals would result in a reduction goal of 49% below 2005 levels by 2035.

Ultimately, the City will determine the level of reductions that it can and should achieve. The recommended targets provided below are guidance based on consistency with the State's goals.

#### **Recommended Community Targets**

In 2020, the City is expected to meet a State-aligned reduction target through existing efforts and legislation. In 2035, the City would need to reduce 44,270 MT CO<sub>2</sub>e emissions below the 2035 Adjusted BAU scenario to meet the State-aligned target (Table 17 and Figure 7).

Sector	2005	2012	2020	2035
BAU Emissions (MT CO <sub>2</sub> e)	287,025	263,876	260,072	259,792
Adjusted BAU Emissions (MT CO <sub>2</sub> e)	287,025	263,876	229,481	190,653
State-Aligned Target (% change from 2005)			-15%	-49%
State-Aligned Target (% change from 2012)			-8%	-45%
State-Aligned Emissions Goal (MT CO <sub>2</sub> e)			243,971	146,383
Reductions from Adjusted BAU needed to meet the Target (MT $CO_2e$ )			Target Met	44,270

Table 17. State-Aligned Community GHG Reduction Targets

<sup>&</sup>lt;sup>4</sup> In an analysis, the State concluded that a 15% reduction in emissions from 2005 levels by 2020 would be equivalent to achieving 1990 emissions levels.




Figure 7. Community Emissions Inventories, Projections, and Targ

#### **Recommended Municipal Targets**

In 2020, the City will need to reduce its emissions by 268 MT CO<sub>2</sub>e from the Adjusted BAU forecast to achieve a reduction goal consistent with the State (Table 18 and Figure 8). Beyond 2020, the City will need to implement additional measures to meet more stringent long-term reduction goals. By 2035, the City will need to reduce municipal operation emissions by 1,044 MT CO<sub>2</sub>e from a 2035 Adjusted BAU forecast to meet a 49% reduction goal (below 2005 levels).

	2005	2012	2020	2035
BAU Emissions (MT CO <sub>2</sub> e)	2,264	2,291	2,291	2,291
Adjusted BAU Emissions (MT CO <sub>2</sub> e)	2,264	2,291	2,177	2,177
State-Aligned Target (% change from 2005)			-15%	-49%
State-Aligned Target (% change from 2012)			-16%	-50%
State-Aligned Emissions Goal (MT CO <sub>2</sub> e)			1,924	1,155
Reductions from Adjusted BAU needed to meet the Target (MT $CO_2e$ )			253	1,022

Table 18. State-Aligned Municipal GHG Reduction Targets




Figure 8. Municipal Emissions Inventories, Projections, and Targets

### **Conclusions and Next Steps**

This Report presents the City's community and municipal inventories, forecasts, and recommended reduction targets. It is the foundation of the EECAP and provides the City a first look at what will be needed to meet emissions reductions that are aligned with the State and to mitigate the City's impacts on climate change. This Report also helps to guide the City in determining feasible energy efficiency reduction opportunities by detailing energy-related emissions, including electricity and natural gas from Residential and Commercial/Industrial sectors.

The next steps in the EECAP process are to review the information provided in this Report and to determine preliminary GHG reduction targets for the community and municipal operations. The South Bay Cities Council of Governments will also begin to work with the City to identify local and subregional energy efficiency measures that could be implemented to reach the City's emissions targets.

# References

Association of Environmental Professionals. 2013. The California Supplement to the United States Community-Wide Greenhouse Gas (GHG) Emissions Protocol.

California Air Resources Board. 2010. Local Government Operations Protocol For the quantification and reporting of greenhouse gas emissions inventories version 1.1.

California Air Resources Board. 2011. California Greenhouse Gas Emissions Inventory: 2000-2009.

ICLEI 2012. U.S. Community Protocol for Accounting and Reporting of Greenhouse Gas Emissions version 1.0.

Intergovernmental Panel on Climate Change. (IPCC) 1996. Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories.

Intergovernmental Panel on Climate Change. 2007. Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M.Tignor and H.L. Miller (eds.). Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

### **Appendix A: Glossary of Terms**

**Adjusted Business-as-Usual:** A GHG forecast scenario that accounts for known policies and regulations that will affect future emissions. Generally, these are state and federal initiatives that will reduce emissions from the business-as-usual scenario.

Baseline Year: The inventory year used for setting targets and comparing future inventories against.

**Business-as-Usual (BAU):** A GHG forecast scenario used for the estimation of greenhouse gas emissions at a future date based on current technologies and regulatory requirements and in the absence of other reduction strategies.

**Carbon Dioxide Equivalent (CO<sub>2</sub>e):** This is a common unit for normalizing greenhouse gases with different levels of heat trapping potential. For carbon dioxide itself, emissions in tons of  $CO_2$  and tons of  $CO_2$ e are the same, whereas one ton of nitrous oxide emissions equates to 298 tons of  $CO_2$ e and one ton of methane equates to 25 tons of  $CO_2$ e. The values are based on the gases' global warming potentials.

**Community Inventory**: GHG emissions that result from the activities by residents and businesses in the city. An inventory reports emissions that occur over a single calendar year.

**Emissions Factor:** A coefficient used to convert activity data into greenhouse gas emissions. The factor is a measure of the greenhouse gas intensity of an activity, such as the amount of  $CO_2$  in one kilowatt-hour of electricity.

**Global Warming Potential (GWP):** The relative effectiveness of a molecule of a greenhouse gas at trapping heat compared with one molecule of CO<sub>2</sub>.

**Metric Ton (MT):** Common international measurement for the quantity of greenhouse gas emissions. A metric ton is equal to 2205 lbs. or 1.1 short tons.

**Municipal Inventory:** GHG emissions that result from the activities performed as part of the government operations in the city and are a subset of the community inventory. An inventory reports emissions that occur over a single calendar year.

**Reduction targets:** GHG emissions levels not to be exceeded by a specific date. Reduction targets are often informed by state recommendations and different targets may be established for different years.

**Sector**: A subset of the emissions inventory classified by a logical grouping such as economic or municipal-specific category.

### **Appendix B: Methodology**

This appendix provides a detailed description of the data sources, emission factors, policies, and assumptions used to develop the greenhouse gas (GHG) emissions inventories, forecasts under a business-as-usual (BAU) scenario, forecasts under an Adjusted BAU scenario, and the recommended GHG reduction targets.

#### Protocols

The GHG inventories for 2005, 2007, 2010, and 2012 were calculated using tools and guidance documents developed or supported by government agencies. Calculation protocols have been developed to ensure consistency among community and municipal inventories. Specifically, the U.S. Community Protocol for Accounting and Reporting of Greenhouse Gas Emissions (Community Protocol) (ICLEI 2012) and the California Supplement (AEP 2013) were used for the community inventories and the Local Government Operations Protocol (LGOP) was used for the municipal inventories (CARB 2010). These protocols often have multiple calculation methods for a single emission source depending on the data available. There are two broad approaches for calculating emissions: "bottom-up" and "top-down." A bottom-up approach relies on end-use data, such as the city-level electricity usage. A top-down approach relies on aggregated data that is allocated to the city based on population, employment, or other relevant indicator. Bottom-up calculations were performed whenever possible to provide the most detailed and likely accurate picture of emissions within a jurisdiction; however, when detailed data were not available, other appropriate methods were used and are described in this appendix. Data were also calculated and managed to best fit the GHG inventory and planning software tool used for this project, called ClearPath. ClearPath was developed by the Statewide Energy Efficiency Collaborative (SEEC) which is a partnership between several statewide agencies, utilities, and non-profits to assist cities and counties in climate mitigation planning. ClearPath is further described at californiaseec.org. In addition, a South Bay Cities Council of Governments (SBCCOG) User's Guide is being developed as part of this project to help cities and SBCCOG to maintain the data and provide for consistent reporting of emissions over time.

#### **Global Warming Potential Factors**

The inventories include the three GHGs most relevant to community and municipal emissions:  $CO_2$ ,  $CH_4$ , and  $N_2O$ . Each GHG differs in its ability to absorb heat in the atmosphere based on their molecular properties and expected lifetime in the atmosphere, and it is useful to describe emissions in one unit of measurement. That unit of measurement is a  $CO_2$ -equivalent, or  $CO_2e$  and Global Warming Potential (GWP) factors are used to standardize emissions from various GHGs. GWP factors, developed by the Intergovernmental Panel on Climate Change (IPCC), represent the heat-trapping ability of each GHG relative to that of  $CO_2$ . For example, the GWP factor of  $CH_4$  is 25 because one metric ton (MT) of  $CH_4$  has 25 times the heat-trapping capacity as one MT  $CO_2$  (over a 100-year period). IPCC periodically updates the GWP factors of GHGs based on new science and updated background mixing ratios of  $CO_2$ .  $CO_2$  always has a GWP factor of 1 and the other GHGs are calculated relative to  $CO_2$ . The California Air Resources Board (CARB) recently updated their GWP factors to align with the IPCC's Fourth Assessment Report, as shown in Table B-1. GWP factors are unitless. Emissions in the inventories are reported in units of  $CO_2e$ .

	CO2	CH <sub>4</sub>	N₂O
GWP	1	25	298

#### **Table B-1. Global Warming Potentials**

Source: IPCC Fourth Assessment Report, 2007.

#### Activity Data

Activity data is the end-use consumption amount of a sector, such as kilowatt hours of electricity, therms of natural gas, and vehicle miles traveled for on-road transportation. In estimating the City's historic GHG emissions, activity data at the City level were obtained when possible (a "bottom-up" approach). When not available, other data sources were used, generally at the county level (a "top-down" approach). Municipal data for 2005 and 2007 were obtained from the City's previous inventory report. Other data were provided by the sources as identified Table B-2.

Data	Data Source	Notes
Community Electricity	Southern California Edison	
Municipal Electricity	Southern California Edison	Maintained by SBCCOG
Community Natural Gas	Southern California Gas Company	
Municipal Natural Gas	Southern California Gas Company	
Community Water	California Water Service Company	
Vehicle Miles Traveled	Southern California Association of Governments (SCAG)	Origin-destination approach, described below
Demographic Data	SCAG	
Vehicle Fleet	City	
Employee Commute	City	
Off-Road Emissions	OFFROAD Model	County-level data
Waste	CalRecycle	

#### **Origin-Destination VMT**

For the community inventory, activity data (vehicle miles traveled) were based on an origin-destination approach used by the State in developing emissions target for metropolitan planning organizations under SB 375. This approach has also been the typical approach used in estimating emission within a city. This approach accounts for:

- Half of the emissions where one endpoint is in the City, for example either the origin or destination of the trip.
- All of the emissions where the trip begins and ends within the City.
- None of the emissions that are "pass-through"; that is, a trip passes through the City but does not begin or end within its boundary.

This approach is used to account for trips or portions of trips that the city may have some control over.

## **Community Activity Data**

Community activity data are shown in Table B-3, except for off-road emissions, which are shown in Table B-4 for Los Angeles County.

Sector	2005	2007	2010	2012	% Change 2005 to 2012						
On-road Transportation	On-road Transportation										
Total Vehicle Miles Traveled	287,926,109	281,650,326	274,693,126	276,102,859	-4.1%						
Residential Energy											
Electricity (kWh)	126,988,678	128,757,878	126,318,979	127,691,251	0.6%						
Natural Gas (therms)	9,471,187	10,064,521	9,495,769	8,518,368	-10.1%						
Commercial/Industrial Energy											
Electricity (kWh)	54,342,771	57,319,790	62,520,704	64,563,386	18.8%						
Natural Gas (therms)	727,360	697,781	917,031	876,032	20.4%						
Solid Waste											
Landfilled (tons)	33,150	31,592	25,285	16,815	-49.3%						
ADC (tons) <sup>1</sup>	3,168	1,855	382	112	-96.5%						
Water and Wastewater											
Water (Million Gallons)	3,424	3,424	3,040	3,165	-7.6%						
Wastewater (City portion of countywide residents)	0.43%	0.43%	0.42%	0.42%	-2.1%						
Off-road Sources <sup>2</sup> (% of LA County emis	sions attributed	to the City)									
Lawn & Garden (% Households)	0.49%	0.48%	0.47%	0.47%	-3.2%						
Construction (% Building permits)	0.05%	0.09%	0.51%	0.10%	114.4%						
Industrial (% Manufacturing jobs)	0.03%	0.03%	0.03%	0.03%	5.0%						
Light Commercial (% Other jobs)	0.15%	0.16%	0.16%	0.16%	7.8%						
Recreation (Population weighted by income)	0.93%	0.90%	0.87%	0.85%	-8.5%						
Agriculture (% Ag. Jobs)	0.00%	0.00%	0.00%	0.02%							

Table B-3. Activity Data used in 2005, 2007, 2010, and 2012 Community Inventories

1 ADC is Alternative Daily Cover, which is green waste (grass, leaves, and branches) that is used to cover landfill emissions. They are reported separately by CalRecycle and therefore shown separately here.

2 Off-road emissions are available at the county level through CARB's OFFROAD model. Emissions attributable to the City were derived using indicator data related to the off-road source. For example, the percentage of households in the City compared to the county was used to attribute the same percentage of lawn & garden equipment emissions to the City. See Appendix B for more methodology details.

Off-road Class	GHG Type	2005 (MT CO₂e /yr)	2007 (MT CO₂e /yr)	2010 (MT CO₂e /yr)	2012 (MT CO₂e /yr)
	CO <sub>2</sub>	921.79	910.27	893.24	882.09
Agricultural Equipment	$CH_4$	0.19	0.17	0.14	0.12
	N <sub>2</sub> O	0.01	0.01	0.01	0.01
	CO <sub>2</sub>	268,646.23	277,541.76	290,911.26	299,875.79
Construction and Mining Equipment	$CH_4$	34.12	31.44	28.24	26.28
Equipment	N <sub>2</sub> O	0.22	0.24	0.25	0.26
	CO <sub>2</sub>	8,099.90	8,562.29	9,255.58	9,870.65
Industrial Equipment	CH <sub>4</sub>	7.16	6.2	4.46	3.89
	N <sub>2</sub> O	0.69	0.63	0.56	0.55
	CO <sub>2</sub>	2,581.13	2,737.30	2,968.71	3,215.02
Lawn and Garden Equipment	$CH_4$	4.98	4.87	4.76	4.96
	N <sub>2</sub> O	2.01	2.01	2.01	2.13
	CO <sub>2</sub>	5,300.36	5,572.36	5,979.92	6,387.77
Light Commercial Equipment	$CH_4$	2.83	2.54	2.18	2.05
	N <sub>2</sub> O	0.91	0.97	1.02	1.07
	CO <sub>2</sub>	286.54	309.8	343.68	369.04
Recreational Equipment	CH <sub>4</sub>	2.14	2.32	2.58	2.77
	N <sub>2</sub> O	0.52	0.57	0.64	0.68

Table B-4. Emissions from Off-road Categories for Los Angeles County

## **Municipal Activity Data**

Municipal activity data are shown in Table B-5.

## **Employee Commute**

Data for Employee Commute in ClearPath are entered as gasoline or diesel. Annual vehicle miles traveled is entered as is the percent of miles traveled by passenger cars, light trucks, and heavy trucks. City staff personnel have not changed significantly since the previous employee commute survey was conducted (see the 2009 City of Rancho Palos Verdes Municipal Greenhouse Gas Emissions Inventory Report for details). Therefore, travel patterns were assumed to be similar and were used for the 2010 and 2012 inventories.

Sector	2005	2007	2010	2012	% Change 2005 to 2012					
Buildings & Facilities										
Electricity (kWh)	603,832	751,034	408,304	656,297	9%					
Natural Gas (therms)	2,391	3,507	261	923	-61%					
Outdoor Lights										
City-Owned (kWh)	114,813	107,983	83,253	102,990	-10%					
SCE-Owned (kWh)	1,173,733	1,173,903	1,175,016	1,167,116	-1%					
Fleet & Equipment										
City-Owned Fleet										
Gasoline (gallons) <sup>1</sup>	3,316	2,738	2,738	2,738	-17%					
Contracted Fleet	· · · ·									
Gasoline (gallons)	24,493	26,617	10,411	16,576	-32%					
Diesel (gallons) <sup>1,23</sup>	101,226	90,259	90,259	90,259	-11%					
LPG (gallons)	16,345	17,761	41,707	48,422	196%					
CNG (standard cubic feet)	1,353,335	1,470,499	788,785	783,137	-42%					
Employee Commute <sup>1, 3</sup>	·			<u> </u>	<u>.</u>					
Gasoline (vehicle miles traveled)	405,885	446,565	397,550	397,550	-2%					
Diesel (vehicle miles traveled)	0	868	14,636	14,636						
# Full Time Employee Equivalents	60	66	66	66	10%					
Solid Waste <sup>1</sup>										
Generated Waste (tons)	24	26	26	26	9%					
Water Delivery										
Electricity (kWh)	100,628	121,593	87,088	138,731	38%					

#### Table B-5. Activity Data used in 2005, 2007, 2010, and 2012 Municipal Inventories

1 Data for 2010 and 2012 were not available; therefore, activity data from 2007 data was used as a proxy.

2 ULSD gallons were used for Diesel.

3 Employee Commute survey conducted in 2014 and adjusted based on the number of employees in 2010 and 2012.

# **Emission Factors**

Emissions factors are used to convert activity data to GHG emissions. An emission factor is defined as the average emission rate of a given GHG for a given source, relative to units of activity. By definition, an emission factor is related to activity data. The emission factors used in the inventories are described by sector below.

## Electricity

California utilities report the average  $CO_2$  content per output of electricity on an intermittent basis. The  $CO_2$ -intensity of electricity varies by utility and year, due to changes in supply, renewable generation, and other factors. The community and municipal operations use electricity provided by SCE except for embedded energy in water, which travels throughout the state and therefore utilizes electricity from multiple utilities (and are shown under the Water Sector).

## Southern California Edison

SCE reported  $CO_2$  factors for 2005 and 2007 through the Climate Registry, and a  $CO_2$ e factor for 2012 in their 2012 Corporate Responsibility & Sustainability Report. When an emission factor is unknown for a certain year, it is standard to use the most recently-reported historic factor until (and if) there is an updated factor. There is no published SCE emission factor for 2010; therefore the factor for 2007 was used for SCE electricity-related emissions calculations in 2010 (Table B-6).

Year	CO2	CH₄	N <sub>2</sub> O	Proxy Year	Data Source
2005	665.72	0.03	0.011	NA	CO <sub>2</sub> : Climate Registry. CH <sub>4</sub> and N <sub>2</sub> O: U.S. Community Protocol
2007	630.89	0.029	0.010	NA	$CO_2$ : Climate Registry. CH <sub>4</sub> and N <sub>2</sub> O: U.S. Community Protocol
2010	630.89	0.029	0.010	2007	$CO_2$ : Climate Registry. CH <sub>4</sub> and N <sub>2</sub> O: U.S. Community Protocol
2012	705 <sup>1</sup>	NA	NA	NA	2012 Corporate Responsibility & Sustainability Report

NA: Not Applicable.

1 The 2012 factor was reported as  $CO_2e$ ; therefore, there are no  $CH_4$  and  $N_2O$  factors.

## **Natural Gas Combustion**

Emission factors for natural gas do not vary greatly over time or by supplier. Therefore, emission factors are U.S. averages as listed in the Community Protocol and are applied for all years (Table B-7).

	CO2	CH₄	N <sub>2</sub> O	Data Source
kg /MMBtu	53.02	0.005	0.0001	U.S. Community Protocol

## **Transportation and Mobile Sources**

#### EMFAC Model

CO<sub>2</sub> emission factors for transportation and mobile sources are calculated using the State-developed Emissions Factor (EMFAC) model, which can be downloaded at <u>http://www.arb.ca.gov/emfac/</u>. Emissions are available at the county level and emission factors were developed and applied to vehicle miles traveled specific to each inventory year. Data are aggregated as annual emissions for all vehicle model years and speeds, but separated by vehicle category. Vehicle categories include light-duty autos,

light-duty trucks, medium-duty vehicles, heavy-duty trucks, and motorcycles.<sup>1</sup> These categorizations are used to develop an emissions factor for gasoline and diesel vehicles. Emission factors were developed using total CO<sub>2</sub> exhaust, which includes emissions from vehicles in motion, idling, and ignition. While emissions from idling and ignitions are not directly related to mileage, they were included so that reductions from measures that may decrease idling could be accounted for in future inventories.

## **On-Road Transportation**

Emissions were converted to emission factors as grams of  $CO_2$  per mile for gasoline and diesel vehicle using EMFAC and a 3-step process (for each inventory year):

- 1. Calculate the vehicle-class average fuel efficiency (miles/gallon) using EMFAC vehicle miles traveled and gallons of fuel consumed for Los Angeles County;
- 2. Calculate the vehicle-class average CO<sub>2</sub> emission factor using EMFAC CO<sub>2</sub> emissions<sup>2</sup> and gallons of fuel consumed for Los Angeles County;
- 3. Calculate the average grams CO<sub>2</sub>/mile traveled factor weighted by vehicle class miles traveled for Los Angeles County.

EMFAC does not provide emissions for  $CH_4$  and  $N_2O$ ; therefore, factors from the Community Protocol were used (Table B-8).

		n Road Aver grams/mile)	•	Diesel On Road Average Factor (grams/mile)			
	CO <sub>2</sub> CH <sub>4</sub> N <sub>2</sub> O			CO2	CH₄	N₂O	
2005	466.062	0.030	0.034	1329.797	0.001	0.001	
2007	464.019	0.028	0.029	1331.634	0.001	0.001	
2010	458.638	0.028	0.029	1280.045	0.001	0.001	
2012	442.657	0.028	0.029	1302.653	0.001	0.001	

**Table B-8. Fleet-Average Emission Factors** 

## **Employee Commute**

Emissions from employee commute in the municipal operations are calculated using annual vehicle miles traveled for gasoline and diesel.  $CO_2$  emissions are estimated using a default emission factor of 8.78 and 10.21 kg/gallon for gasoline and diesel, respectively<sup>3</sup> and fuel economy, which is based on EMFAC outputs for each inventory year and vehicle class. Vehicle miles traveled are converted to  $CH_4$  and  $N_2O$  emissions using emission factors from the Community Protocol. Table B-9 shows the miles per

<sup>&</sup>lt;sup>1</sup> Vehicle categories may use either EMFAC2007 or EMFAC2011 categorizations and result in the same data for the purposes of these inventories; EMFAC2007 categories were used here EMFAC2011 further disaggregates medium heavy-duty vehicles and heavy heavy-duty vehicles into 29 vehicle categories. This level of detail is not needed for these inventories. More information on vehicle categories is available at <a href="http://www.arb.ca.gov/msei/vehicle-categories.xlsx">http://www.arb.ca.gov/msei/vehicle-categories.xlsx</a>.

<sup>&</sup>lt;sup>2</sup> For 2010 and 2012, the emissions accounting for the effects of existing policies (Pavley and Low Carbon Fuel Standard) were used. These standards did not exist in 2005 and 2007.

<sup>&</sup>lt;sup>3</sup> Information from ClearPath developers e-mail dated June 19, 2014.

gallon and grams (CH<sub>4</sub> and N<sub>2</sub>O) per mile used to estimate emissions from employee commute by vehicle class.

#### **Vehicle Fleet**

Vehicle fleet consists of City-owned and contracted vehicles used to perform City services. Vehicle Fleet requires input of gallons of fuel used by fuel type to estimate  $CO_2$  emissions. Vehicle miles traveled are used to estimate  $CH_4$  and  $N_2O$ . The factors used for the City are shown in Table B-9.

		2005	2007	2010	2012						
Gasoline	Gasoline										
	MPG	21.700	21.875	22.027	22.064						
Passenger Vehicle	g CH₄/mi	0.030	0.028	0.028	0.028						
	g N₂O/mi	0.034	0.029	0.029	0.029						
	MPG	16.575	16.666	16.795	16.823						
Light Truck	g CH₄/mi	0.035	0.031	0.031	0.031						
	g N₂O/mi	0.049	0.043	0.043	0.043						
	MPG	12.754	12.806	12.854	12.856						
Heavy Truck	g CH₄/mi	0.033	0.033	0.033	0.033						
	g N₂O/mi	0.013	0.013	0.013	0.013						
Diesel											
	MPG	27.558	27.662	29.006	29.889						
Passenger Vehicle	g CH₄/mi	0.001	0.001	0.001	0.001						
	g N₂O/mi	0.001	0.001	0.001	0.001						
	MPG	27.032	27.251	27.705	28.498						
Light Truck	g CH₄/mi	0.001	0.001	0.001	0.001						
	g N₂O/mi	0.001	0.001	0.001	0.001						
	MPG	17.343	17.588	18.797	18.858						
Heavy Truck	g CH₄/mi	0.005	0.005	0.005	0.005						
	g N₂O/mi	0.005	0.005	0.005	0.005						

Note: MPG is miles per gallon and is derived from EMFAC at the county level.  $CH_4$  and  $N_2O$  emission factors are from the Community Protocol; Passenger Vehicle and Light Truck emission factors have data for 2005 and later; Heavy Truck only have 2010 data.

## **Off-Road**

Off-road emissions include emissions from agriculture, construction, industrial, lawn and garden, light commercial, and recreational equipment. Annual emissions of  $CO_2$ ,  $CH_4$ , and  $N_2O$  are available at the county level from the State's OFFROAD model. To estimate values for each city, relevant indicator data are used to estimate the proportion of county-level emissions attributable to the city. Table B-10 lists

the indicator used to estimate the City's portion of emissions for each category and Table B-11 shows City-specific data. City- and county-level indicator data were obtained from SCAG.

Category	Indicator
Agriculture Equipment	Agriculture Jobs
Construction Equipment	Building Permits Issued
Industrial Equipment	Manufacturing Jobs
Lawn and Garden Equipment	Households
Light Commercial Equipment	Non- Manufacturing or Agriculture Jobs
Recreational Equipment	Population, Weighted by Median Income

Table B-10. Off-road Emissions Indicators

		Ag. Jobs	Building Permits	Mfg. Jobs	Households	Other Jobs <sup>1</sup>	Population	Income (\$)
	City	0	12	124	15,417	6,067	42,457	104,347
2005	County	13,562	25,623	461,099	3,178,736	4,045,922	9,816,200	48,606
	%	0.00%	0.05%	0.03%	0.49%	0.15%		0.93%
	City	0	19	132	15,480	6,444	41,728	108,296
2007	County	13,562	20,303	461,099	3,224,053	4,045,922	9,780,800	51,439
	%	0.00%	0.09%	0.03%	0.48%	0.16%		0.90%
	City	0	38	104	16,179	6,028	41,654	114,500
2010	County	10,598	7,466	362,157	3,454,093	3,758,244	9,818,605	56,000
	%	0.00%	0.51%	0.03%	0.47%	0.16%		0.87%
	City	2	19	104	16,218	6,192	41,897	108,046
2012	County	10,798	18,926	369,005	3,454,093	3,829,313	9,889,632	53,880
	%	0.02%	0.10%	0.03%	0.47%	0.16%		0.85%

#### Table B-11. Off-road Emissions Indicator Data

Note: Some percentages may appear off due to rounding. Ag. = Agriculture. Mfg. = Manufacturing.

1 Other indicates non-manufacturing and non-agricultural.

#### Water

Emissions from water are indirect. Water requires energy to move from its source to final treatment and the energy for most of these processes is not captured in local utility data (i.e., the portion that is used in a home or business and therefore contained in the owner's utility bill). This portion is termed the "embedded energy" in water and particularly for southern California, the energy embedded in water is high and should be accounted for in a community inventory. The California Energy Commission (CEC) developed a report, titled <u>Refining Estimates for Water-Related Energy Use in California</u>, which estimates the energy required to supply, convey, distribute, and treat water in northern and southern California. Recycled water is less energy-intensive because it does not require the supply and conveyance energy. Outdoor water infiltrates into the ground and therefore does not have the

wastewater energy treatment component. Therefore, the emission factors are adjusted to account for the proportion of recycled and outdoor water. The amount of water used for indoor or outdoor use was not available at the City level; however, the 2010 Los Angeles Department of Water & Power, Urban Water Management Plan states that 61% of water is for indoor use for the City of Los Angeles. The water usage is assumed to be similar for the South Bay sub-region. Therefore, the embedded energy in a million gallon (MG) of water in the City is estimated in Table B-12 using the CEC report and estimated indoor vs. outdoor water usage in the region.

	Conventional <sup>1</sup> (kWh/MG)	Recycled (kWh/MG)
Supply and Convey	9,727	
Treatment	111	111
Distribution	1,272	1,272
Wastewater Treatment	1,911	1,911
Total	13,022	3,294
South Bay Factor	12,275.71	2,548.71

Table B-12. Energy Embedded in Water

1 From CEC's 2006 Refining Estimates for Water-Related Energy Use in California, for Indoor water use in southern California.

#### **Statewide Average Electricity**

For energy embedded in water, a statewide average emission factor is applied because water in the South Bay sub-region is supplied from various regions in the State (Table B-13). Similar to SCE data, statewide emission factors are not available for each inventory year. For 2010 and 2012, the 2009 statewide emission factors were used as the proxy year.

Year	CO2	CH₄	N <sub>2</sub> O	Proxy Year	Data Source
2005	948.28	0.03	0.011	NA	U.S. Community Protocol
2007	919.64	0.029	0.010	NA	U.S. Community Protocol
2010	658.68	0.029	0.006	2009	U.S. Community Protocol
2012	658.68	0.029	0.006	2009	U.S. Community Protocol

Table B-13. California Statewide Electricity Emission Factors

NA: Not Applicable.

## Wastewater

The emissions for wastewater include the  $CH_4$  and  $N_2O$  emissions from processing which consist of three sources: **stationary**, **process**, and **fugitive** emissions.

**Stationary** emissions are derived from combustion of digester gas at a centralized treatment facility. The City is served by the Los Angeles County Sanitation District's Joint Water Pollution Control Plant (JWPCP). JWPCP is a centralized treatment facility that uses an anaerobic digester process and does not employ a formal nitrification/denitrification (N/DN) system. Detailed information regarding the amount of digester gas produces was not available, so an alternative method using City population information

264<sup>d</sup>

41.1<sup>d</sup>

was used. Default factors from the Community Protocol were applied to estimate  $CH_4$  and  $N_2O$  emissions for stationary emissions. Although  $CO_2$  emissions are also produced, the fuel source is considered a biofuel, and the resulting  $CO_2$  emissions are considered "biogenic" and are not reported<sup>4</sup>.

**Process** emissions include N<sub>2</sub>O emissions as a result of N/DN processes at the treatment facility. All wastewater facilities have emissions from N/DN—some facilities have a formal N/DN process, which would result in greater N/DN emissions, but for the JWPCP, N/DN emissions are solely a result of natural processes. The recommended approach to estimating these emissions is through the population served and default factors listed in the Community Protocol. In an advanced, centralized treatment facility, stationary and process emissions are relatively small compared to fugitive emissions. The Community Protocol, and likewise ClearPath, recommends multiplying the population-derived emissions by 1.25 to account for commercial and industrial discharges to the system. Regions without any commercial and industrial sources should use a factor of 1.0. Because the City is largely residential, a factor of 1.0 was applied to these emissions.

**Fugitive** emissions occur from inflow (septic systems) and effluent discharge. JWPCP reports facilitywide effluent, and effluent nitrogen content, which are factors used in estimating fugitive emissions (Table B-14). The City's portion was determined by estimating the proportion of the population served by JWPCP. The ClearPath tool requires the daily N load in kg N per day. This is calculated using the factors listed in Table B-14 and the Community Protocol Equation WW.12:

Daily N Load for the City (kg N/day) = Effluent X Effluent Nitrogen Content X gallons/liter X City Population/Service Population,

Where Effluent is the facility-wide discharge in millions of gallons per day (MGD), Effluent Nitrogen Content is the average nitrogen content per volume (mg/L), and gallons/liter is a conversion factor (3.79). The Daily N Load entered into ClearPath was adjusted by a factor of 0.5 to account for the difference in emission factors for direct ocean discharge and stream/river discharge. In ClearPath, ocean discharge is not an option; however, the emissions are estimated to be ½ of those from discharge to a stream or river (see Community Protocol Appendix F). Therefore, the Daily N Load was adjusted by 0.5 to account for this difference.

in Wastewater Fugitive Emissions					
	2005	2007	2010	2012	

403<sup>ª</sup>

296<sup>b</sup>

36.7<sup>b</sup>

237<sup>c</sup>

39.7<sup>e</sup>

Table B-14. Los Angeles County Joint Water Pollution Control Plant Data Used	
in Wastewater Fugitive Emissions	

Effluent Nitrogen content (mg/L)	
a Default assumption based on influent.	

b 2008 annual report data.

Effluent (MGD)

c 2011 annual report data.

d 2013 annual report data.

e Based on communication with Los Angeles County Sanitation District for 2009.

<sup>&</sup>lt;sup>4</sup> Emissions from digester gas combustion are automatically calculated in ClearPath when population is entered.

#### Solid Waste

Emissions from solid waste are primarily in the form of fugitive emissions of methane from decomposition. Emission factors are derived from the Community Protocol, based on the type of waste disposed. The State conducts a Waste Characterization Study (Study) every 4 to 6 years to determine the amount of waste attributable to each waste type. The Study is conducted at the State level by economic sector; therefore, community-level characterizations are not available. For the community inventory, the overall composition of California's disposed waste stream was used to convert total tons into waste types (Table B-15). For the municipal inventory, the characterization for public administration was used (Table B-15). In addition to community-generated waste, some diverted green waste is used as landfill cover rather than importing landfill cover from other regions. This green waste is known as alternative daily cover (ADC) and is reported by CalRecycle for each community. The ADC characterization was determined through communication with the developers of ClearPath and does not vary by year or community. The emission factor to determine methane generation varies if the landfill operates a methane flare or generates electricity from methane capture. The Community Protocol recommends using an average factor of 75% recovery from landfill gas, although some landfills with have much higher gas recovery systems, and other landfills do not have any. Carbon dioxide generated by decomposition of waste in landfills is not considered anthropogenic because it would be produced through the natural decomposition process regardless of its disposition in the landfill. Nitrous oxide is not a by-product of decomposition and therefore no fugitive emissions of nitrous oxide are anticipated from this source. The waste characterizations and emission factors used to estimate emissions from solid waste are provided in Table B-15. The "Category in in the 2004 and 2008 Studies" detail which Study categories make up the ClearPath Category.

ClearPath Category	Category in 2004 and 2008 Studies	Alternative Daily Cover <sup>1</sup>	2004 Study <sup>2</sup>	2008 Study <sup>3</sup>	Public Administration	Emission Factor <sup>1</sup>
Newspaper	Newspaper	0%	2.2%	1.3%	5.5%	0.043
Office Paper	White/Colored Ledger Paper + Other Office Paper + Other Miscellaneous Paper	0%	5.4%	4.9%	13%	0.203
Cardboard	Uncoated Corrugated Cardboard + Paper Bags	0%	6.7%	5.2%	5.1%	0.120
Magazine/ Third Class Mail	Magazines and Catalogs + Remainder/ Composite Paper	0%	6.5%	5.9%	15.4%	0.049
Food Scraps	Food	0%	14.6%	15.5%	9.8%	0.078
Grass	Leaves and Grass	30%	2.1%	1.9%	8.05%	0.038
Leaves	Leaves and Grass	40%	2.1%	1.9%	8.05%	0.013
Lumber	Branches and Stumps + Prunings and Trimmings	0%	2.6%	3.3%	0.1%	0.062
Branches	Lumber	30%	9.6%	14.5%	5%	0.062

Table B-15. W	/aste Characterization a	nd Emission Factors	for Solid Waste
---------------	--------------------------	---------------------	-----------------

1 Breakdown from ClearPath Developers via e-mail dated June 19, 2014. Used for all inventory years.

2 2004 Waste Characterization Study for California, Overall Waste Stream. Used for 2005 inventory. Does not total 100% as not all waste is organic.

3 2008 Waste Characterization Study for California, Overall Waste Stream Used for 2007, 2010, 2012 inventories. Does not total 100% as not all waste is organic.

## Forecasts

The forecasts are an estimate of what emissions in the City may be in 2020 and 2035. The forecasts were developed using standard methodologies under two scenarios: Business-as-Usual (BAU) and Adjusted BAU.

## **Business-as-Usual Forecasts**

The BAU scenario uses current (2012) consumption patterns and predicted growth in the City in the absence of state and federal legislation that would reduce future emissions. The growth assumptions are those estimated by SCAG in their 2012 Regional Transportation Plan and are applied to emissions sectors based on their relevance. For example, future Residential Energy emissions were developed using current energy use per household (from the 2012 inventory) and the anticipated number of households in the future. Table B-16 shows the growth factors used to project emissions in the City.

#### Table B-16. Emissions Sectors and Demographic Growth Indicators

Sector	Demographic Indicator	
Residential Energy	Households	
Commercial/ Industrial Energy	Jobs	
Solid Waste, Water, Wastewater, Aviation, Off-Road Sources	Service Population (Population + Jobs)	
Transportation	Vehicle Miles Traveled, modeled by SCAG	
Municipal Jobs	Municipal Emissions <sup>1</sup>	

SCAG: Southern California Association of Governments

1 The number of jobs in the City is used as an indicator for all municipal operation emissions except Aviation, which is forecast consistent with the community forecast (by change in service population).

## **Adjusted Business-as-Usual Forecasts**

The Adjusted BAU scenario also uses growth estimates for the City, also accounts for legislation that will reduce emissions in the future, regardless of City actions. Table B-17 summarizes the legislation that will reduce the City's emissions in the future and which sectors the legislation applies to.

Legislation	Description	Emissions Sector Affected
Low Carbon Fuel Standard	Reduce carbon intensity of transportation fuels 10% by 2020.	On-road Transportation, Employee Commute, Vehicle Fleet
AB 1493 and Advanced Clean Cars	Implement GHG standards for passenger vehicles, implement zero-emission vehicle program, support clean fuels outlet regulation.	On-road Transportation
California Building Code Title 24	Improved energy efficiency standards for new residential and non-residential construction.	Residential Energy, Non-residential Energy
Renewable Portfolio Standard <sup>1</sup>	Provide 33% of electricity from renewable sources by 2020.	Water
Senate Bill X7-7	Reduce urban per capita water consumption 20% by 2020.	Water

1 Potential GHG reductions from this legislation were not applied to the electricity in SCE's service territory due to the uncertainty in SCE's generation sources after the closure of the San Onofre Nuclear Generating Station.

#### Low Carbon Fuel Standard, AB 1493, and Advanced Clean Cars

Changes in on-road emissions in Los Angeles County were modeled using EMFAC, which models both the emissions with and without Low Carbon Fuel Standard and Pavley I. Additional modeling was conducted to estimate the change in emissions due to Advanced Clean Cars. The rate of reductions from on-road transportation measures through 2020 was assumed to be 0.0344% per year for gasoline and 0.0106% per year for diesel. After 2020, the rate of reductions was assumed to be 0.03452% per year for gasoline and 0.0251% per year for diesel.

### **California Building Code Title 24**

Title 24 updates will raise the minimum energy efficiency standards for new buildings, thereby decreasing the expected energy consumption of future development in the City. Under the adjusted BAU scenario, it was assumed that the 2013 Title 24 standards that went into effect in 2014 will make new residential and non-residential buildings more efficient than they would be under the 2008 Title 24 standards for new residential buildings. The energy savings were estimated using analyses developed by the California Energy Commission and the applied to the expected new development in the City to 2020 and 2035. The rate of reductions was applied to the City's 2012 energy use (kWh or therms) per household (for Residential energy) or per job (for Commercial energy). Savings were applied to new development anticipated in the City. Detailed energy savings assumptions are below.

### Residential

Residential electricity is estimated to be 32.6% lower under the new standards.<sup>5</sup> This percentage savings is relative to heating, cooling, lighting and water heating only and do not include other appliances, outdoor lighting that is not attached to buildings, plug loads, or other energy uses. Electricity consumption due to heating, cooling, lighting, and water heating accounts for 34% of total household electricity use.<sup>6</sup> Therefore, the percentage of total residential electricity that will be reduced as a result of the 2013 Title 24 standards is 11.1%.

Residential natural gas savings were estimated 5.8% lower under the new standards. Again, this percentage savings pertains only to the energy sources affected by Title 24 Standards. Natural gas consumption due to space and water heating accounts for 86% of total household natural gas use.<sup>7</sup> Therefore, the percentage of total residential natural gas that will be reduced as a result of the 2013 Title 24 standards is 5.0%.

## Commercial

Commercial Electricity savings were estimated to be 21.8% lower under the new standards. Title 24related measures would impact 77.2% of total electricity use in commercial buildings<sup>8</sup>; therefore, 16.8% reduction in electricity consumption may be expected in new commercial development.

<sup>&</sup>lt;sup>5</sup> CEC Impact Analysis, California's 2013 Building Energy Efficiency Standards, July 2013. CEC-400-2013-008.

<sup>&</sup>lt;sup>6</sup> CEC 2009 California Residential Appliance Saturation Appliance Study, October 2010. CEC-200-2010-004.

<sup>&</sup>lt;sup>7</sup> CEC 2009 California Residential Appliance Saturation Appliance Study, October 2010. CEC-200-2010-004.

<sup>&</sup>lt;sup>8</sup> CEC 2006. California Commercial End-Use Survey. March 2006. CEC-400-2006-005.

Natural gas savings were estimated to be 16.8% under the new standards compared to the previous standards. Heating and cooling account for 69.7% of natural gas consumption in commercial facilities; therefore, 11.7% reduction in natural gas consumption may be expected from 2013 Title 24 standards applied to new commercial development.

## **Renewable Portfolio Standard**

The Renewable Portfolio Standard will be fully implemented in 2020. The level of implementation varies by utility; however, ICLEI estimates that the average statewide level of implementation is 5% per year, compounded annually. As noted in the Report, this reduction is only taken for electricity used in the transport and treatment of water, which moves throughout the State. The reduction is not taken for electricity wholly within SCE's territory.

## Senate Bill X7-7

SB X7-7 will be implemented by individual water districts. The City obtains water from California State Water Service Company's Hermosa/Redondo District and Palos Verdes District. The proportion of water from each district was not available; therefore, the level of implementation of SB X7-7 was estimated using an annualized reduction rate from each of the districts' goals, assuming half of the City's water is derived from each district.

# **Target Setting**

The state-aligned targets are provided to assist the City in determining appropriate emission reduction goals. Recommended targets are based on existing California climate change legislation and State guidance relevant to establishing a GHG reduction target. While State goals are based on a 1990 baseline year, the City's baseline year is 2005. Therefore, the reduction targets are expressed as a percent reduction below 2005 levels. Targets are recommended for 2020 to align with AB 32 and 2035, which is a midpoint between the 2020 goal and the State's long-term 2050 goal. Planning beyond 2035 is considered speculative, as legislation and technology may change significantly before 2050. While it is important for continued reductions well beyond 2035, no local targets are recommended at this time.

Table B-18 provides a summary of the State's goals and the State's guidance to local governments regarding GHG reduction targets. This guidance applies to both municipal operations and communitywide emissions reductions efforts.

	2020	Interim Year Between 2020-2050	2050
State Targets (AB 32 and EO S-3-05)	1990 levels	NA	80% below 1990 levels
State Guidance on Local Government Targets (AB 32) Scoping Plan Recommended Target and Attorney General's Office Guidance	15% below 2005-2008 levels	Demonstrate a trajectory toward statewide 2050 levels (e.g., 49% below 2005 levels by 2035)	NA

 Table B-18. Summary of State Reduction Targets and Guidance on Local Government Targets

 Aligned with State Targets

Table B-19 demonstrates how the local targets are aligned with State targets.

Target Year	Percent below 1990 Emission Levels	Percent below 2005 Emission Levels
2020	0.0%	15.0%
2021	2.7%	17.3%
2022	5.3%	19.5%
2023	8.0%	21.8%
2024	10.7%	24.1%
2025	13.3%	26.3%
2026	16.0%	28.6%
2027	18.7%	30.9%
2028	21.3%	33.1%
2029	24.0%	35.4%
2030	26.7%	37.7%
2031	29.3%	39.9%
2032	32.0%	42.2%
2033	34.7%	44.5%
2034	37.3%	46.7%
2035	40.0%	49.0%

Table B-19. Comparison of 1990 Baseline Targets vs. 2005 Baseline Targets